

Determinants of Anemia in Antenatal Cases: A Cross-sectional Analysis

Dr. Kusum Lata Sharma¹, Dr. Isha Sharma², Dr. Harsh Sharma³, Dr. Puja Dixit⁴,
Dr. Chhaya Tiwari^{5§}

¹Senior Gynecologist, Dr. Kusum Lata Sharma Hospital Bharatpur (Rajasthan) India

²Dentist, Dr. Kusum Lata Sharma Hospital Bharatpur (Rajasthan) India

³Dentist, Dr. Kusum Lata Sharma Hospital Bharatpur (Rajasthan) India

⁴Medical Officers, State Government of Rajasthan India

⁵Gynecologist, Department of Gynecology and Obstetrics, Kailash Hospital, New Delhi, India

Abstract— *Anemia in pregnancy is a major health problem in developing country like India. So this present study was carried out at Dr. Kusum Sharma Hospital Bharatpur (Rajasthan) India, with the aim to find out the determinants of anemia in Antenatal cases attended for delivery. period on pregnancy outcomes. For this study, 100 Antenatal Cases (ANCs) attended for delivery at Dr. Kusum Sharma Hospital Bharatpur (Rajasthan) India were included in this study. These ANCs were interrogated and investigated for hemoglobin estimation. It was found in this study that 63% of proportion of ANCs were having Anemia. Anemia was found to associate with age, residence, education, occupation and parity of ANCs but not with BMI and bad obstetric history of ANCs. Anemia was found significantly more in younger age, less educated and housewives than their counterparts. Likewise ANCs of rural area had more chances of Anemia than those residing in urban areas. And ANCs having either zero parity or parity more than two had more probability to have anemia than their counterparts.*

Keywords: *Antenatal Cases (ANCs), Anemia in pregnancy, Determinants of ANCs*

I. INTRODUCTION

Of the 600,000 deaths from pregnancy related complications world over, anemia is responsible for 40-60% of them.¹ Anemia causes direct as well as indirect maternal deaths from cardiac failure, hemorrhage, infection and pre-eclampsia. Among anemia, iron deficiency anemia is the most common. Nutritional deficiency is the commonest cause for iron deficiency anemia, especially in developing countries like ours.²

Anemia in pregnancy is defined by WHO as a hemoglobin concentration below 11g/dl.³ Although only 15% of pregnant women are anemic in developed countries,⁴ the prevalence of anemia in developing countries is relatively high (33% to 75%).³⁻⁵ According to NFHS-III (2005-2006) prevalence of anemia among pregnant women in India is 58%.⁶

In India, prevalence ranges from 33% to 89%. ICMR district nutrition survey 1999-2000 also reported prevalence of anemia as 84.2% with 13.1% with severe anemia in pregnancy.⁷ Anemia is associated with 22% maternal deaths around the world.⁷ India contributes to about 80% of maternal deaths due to anemia in south Asia.⁸ In India anemia is second most common cause of maternal death, accounting for 20% of the total maternal death.⁹

Iron deficiency is principal cause of anemia.¹⁰ Only 22.3% pregnant women consumed Iron and folic acid tablets for 100 days. (NFHS 2005-06). National Nutrition Monitoring Bureau (NNMB-2003) and RCH surveys have shown that coverage under IFA supplementation was low and even among those who received the tablets, only one-third of them were regularly taking.¹¹

This present study was conducted with the aim to determine the factors associated with anemia in ANCs attended at a private hospital of western Rajasthan.

II. METHODOLOGY

This hospital based cross-sectional observational study was carried out on 100 ANCs attending at Dr. Kusum Sharma Hospital Bharatpur (Rajasthan) India.

Sample size was calculated 100 subjects at 95% confidence limit and 20% relative allowable error assuming 50% prevalence of anemia in pregnant women.

Consecutive ANCs attending at this hospital were included in the study till the sample size i.e. 100 ANCs were achieved. These ANCs were interrogated and data regarding their age, residence, weight and height was noted. Body Mass Index (BMI) of each ANC was calculated as below

$$\text{BMI} = \text{Weight} / \text{Height}^2$$


Information regarding gravidity, parity, and obstetric history was also taken.

Data thus collected were entered in MS Excel 2010 worksheet in the form of master chart. Determinants of anemia was found with Chi-square Test and unpaired 't' test/ANOVA test of significance.

III. RESULTS

In the present study out of 100 ANCs, 63 (i.e. 63%) ANCs were having anemia while only 47 (i.e. 47%) ANCs were not having anemia.(Figure 1)

Figure 1



Regarding bio-social factors, present study also observed that anemia was found to be associated with age and residence but not with BMI. It was revealed that anemia was found more in younger age, <10th standard and house wives than their counterpart groups. Likewise anemia was found more in ANCs from rural areas. (Table 1)

Table 1
Comparison of Bio-social variables of ANC's pregnancy with and without Anemia (N=100)

Bio-social Variables		With Anemia Group (N=63)	Without Anemia (N=47)	P Value	LS
Age (in years)	<20 Years	16	1	0.007*	S
	20-24 Years	24	19		
	25-29 Years	21	23		
	≥30 years	2	4		
Residence	Urban	47	25	0.033*	S
	Rural	16	22		
Education	Illiterate	19	21	0.006*	S
	<10th Standard	35	12		
	>10th Standard	9	14		
Occupation	Housewives	52	28	0.014*	S
	Working	11	19		
BMI	Mean ± SD	20.3±4.2	21.4±5.1	0.218**	NS

*P value with Chi-square

**P value with Unpaired 't' Test

Regarding obstetrical history, anemia was found to be associated with parity but not with bad obstetric history. Anemia was found significantly more in nullipara and ANC's with more than 2 parity than that of parity one or two. (Table 2)

Table 2
Comparison of Variables related to Obstetrical history of ANC's pregnancy with and without Anemia (N=100)

Bio-social Variables		With Anemia Group (N=63)	Without Anemia (N=47)	P Value	LS
Parity	Nullipara	18	7	0.015*	S
	Upto 2	24	31		
	>2	21	9		
Bad OH	Yes	12	9	0.817*	NS
	No	51	38		

*P value with Chi-square

IV. DISCUSSION

In the present study anemia was found in 63% ANC's attended at Dr. Kusum Sharma Hospital Bharatpur. NFHS-III (2005-2006) reported prevalence of anemia among pregnant women in India is 58%.⁶ Another study reports Anemia in 43.38% in ANC's attending at a private hospital of Bareilly district during the month of Jan.2010 to May2010.¹² However higher prevalence of anemia (57.23, 96.5 and 84.9% respectively) had been reported by various other studies.^{13,14,15} This difference is perhaps the present study was limited to the hospital and that to in urban area.

Present study also observed that anemia was found to be associated with age, residence, parity and but not with BMI and bad OH. It was revealed that anemia was found more in younger age group, ANC's from rural areas and in either nullipara or parity more than two. However a cross sectional study was carried out in obstetric and gynecological department OPD of one of the private hospital of Bareilly district during the month of Jan.2010 to May2010.¹² who found more prevalent in women age more than 30 years (80.39%), illiterate (49.53%), working (83.82%). Parity was found to be reported with other authors also.^{14,16}

V. CONCLUSION

ANCs with anemia were found 63%. Anemia was found to be associated with age, education, occupation, parity and residence of ANCs but not with BMI and bad obstetric history. Anemia was found significantly more in younger age, less educated and housewives than their counterparts. Likewise ANCs of rural area had more chances of Anemia than those residing in urban areas. And ANCs having either zero parity or parity more than two had more probability to have anemia than their counterparts.

CONFLICT OF INTEREST

None declared till now.

REFERENCES

- [1] Sharma JB. Nutritional anemia during pregnancy in non industrialized countries. In stud J (ed). Progress in Obstetric and gynaecology vol.15. London, Churchill Livingstone 2003; 103-20.
- [2] Srivastava Aariti, Tondon pabha, Qureshi Sabuhi, Das Vinita. Anemia in pregnancy- A novel regime of intramuscular iron therapy. J Obstet Gynaecol India 2005; 55(3): 237-240.
- [3] World Health Organization (WHO). The prevalence of Anemia in women: a tabulation of available information. Geneva, Switzerland: WHO; 1992. (WHO/MCH/MSM/92.2).
- [4] World Health Organization (WHO). Prevention and Management of severe anemia in pregnancy. Report of a technical working group. Geneva, Switzerland WHO; 1993. (Who/FNE/MSM/93.5).
- [5] Ogunbode. Anemia in pregnancy. In: okonofua F Odunsi, editors. Contemporary obstetrics and gynaecology For Developing countries. Benin city, Nigeria: Women's Health and Action Research Center, 2003. Pp 514-529.
- [6] Kishore J. National Health Programme of India, Century Publication, New Delhi, India, 8th edition, 2010; p 41.
- [7] Stolzfiis, Mullany and Black. Iron Deficiency Anemia."Comparative quantification of health risks: Global and regional burden of disease attributable to selected major risk factors," WHO, Page 163-210, 2004.
- [8] Ezzati M, Lopez AD, Dogers A, Vander HS, Murray C. Selected major risk factors and global and regional burden of disease. Lancet . , 2002; Vol.360, 1347-60
- [9] Iron Deficiency Anaemia: Assessment, Prevention and Control .Geneva, World Health Organization, 2001.
- [10] G M Dhaar, I Robbani. Foundations of Community Medicine, 2nd ed. Elsevier Publishers; 2008
- [11] Abel R, Rajaratnam J, Gnanasekaran VJ and Jayaraman P. Prevalence of anemia and iron deficiency in three trimesters in rural Vellore district, South India. Trop. Doct April 2001; 31(2):86-9
- [12] Kiran Agarwal, Varsha Chaudhary, V.K. Agrawal, Ashok Agarwal, Rajeev Kumar, Mahendra Sharma. Prevalence and Determinants of Anemia in Pregnancy at Private Hospital Of Bareilly District. NJIRM 2011; Vol. 2(4) p29-32
- [13] Vanden Broek NR, Rogerson SJ, Mhango CG, Kambala B, White SA, Molyneux ME. Anemia in pregnancy in southern Malawi: prevalence and risk factors. British Journal of Obstetrics and Gynecology 2000; 107: 445-51.
- [14] Gautam Virender P, Bansal Yogesh, Taneja DK, Saha Renuka. Prevalence of Anemia amongst pregnant women and its socio-demographic associates in a rural area of Delhi. Indian Anemia in pregnancy NJIRM 2011; Vol. 2(4). October-December eISSN: 0975-9840 pISSN: 2230 - 9969 J. Community Medicine 2002; 27 (4): 157-60.
- [15] Toteja GS, Singh P, Dhillon BS, Saxena BN, Ahmed FU, Singh RP et al. Prevalence of anemia among pregnant women and adolescent girls in 16 districts in India. Food Nutr Bull 2006; 27 (4): 311-15
- [16] Abrehet Abriha, Melkie Edris Yesuf and Molla Messele Wassie. Prevalence and associated factors of anemia among pregnant women of Mekelle town: A cross sectional study. BMC research note 1014.7;888